

Harvard Math 129: Algebraic Number Theory

Homework Assignment 3

William Stein

Due: Thursday, March 3, 2005

*The problems have equal point value, and multi-part problems are of the same value. There are **five** problems. In any problem where you use a computer, include in your solution the exact commands you type and their output. You may use any software, including (but not limited to) MAGMA and PARI.*

1. Let $\varphi : R \rightarrow S$ be a homomorphism of (commutative) rings.
 - (a) Prove that if $I \subset S$ is an ideal, then $\varphi^{-1}(I)$ is an ideal of R .
 - (b) Prove moreover that if I is prime, then $\varphi^{-1}(I)$ is also prime.
2. Let \mathcal{O}_K be the ring of integers of a number field. The Zariski topology on the set $X = \text{Spec}(\mathcal{O}_K)$ of all prime ideals of \mathcal{O}_K has closed sets the sets of the form

$$V(I) = \{\mathfrak{p} \in X : \mathfrak{p} \mid I\},$$

where I varies through *all* ideals of \mathcal{O}_K , and $\mathfrak{p} \mid I$ means that $I \subset \mathfrak{p}$.

- (a) Prove that the collection of closed sets of the form $V(I)$ is a topology on X .
- (b) Let Y be the subset of nonzero prime ideals of \mathcal{O}_K , with the induced topology. Use unique factorization of ideals to prove that the closed subsets of Y are exactly the finite subsets of Y along with the set Y .
- (c) Prove that the conclusion of (a) is still true if \mathcal{O}_K is replaced by an order in \mathcal{O}_K , i.e., a subring that has finite index in \mathcal{O}_K as a \mathbb{Z} -module.

3. Explicitly factor the ideals generated by each of 2, 3, and 5 in the ring of integers of $\mathbb{Q}(\sqrt[3]{2})$. (Thus you'll factor 3 separate ideals as products of prime ideals.) You may assume that the ring of integers of $\mathbb{Q}(\sqrt[3]{2})$ is $\mathbb{Z}[\sqrt[3]{2}]$, but do *not* simply use a computer command to do the factorizations.
4. Let $K = \mathbb{Q}(\zeta_{13})$, where ζ_{13} is a primitive 13th root of unity. Note that K has ring of integers $\mathcal{O}_K = \mathbb{Z}[\zeta_{13}]$ (you do not have to prove this).
 - (a) Factor 2, 3, 5, 7, 11, and 13 in the ring of integers \mathcal{O}_K . You may use a computer.
 - (b) For $p \neq 13$, find a conjectural relationship between the number of prime ideal factors of $p\mathcal{O}_K$ and the order of the reduction of p in $(\mathbb{Z}/13\mathbb{Z})^*$.
 - (c) Compute the minimal polynomial $f(x) \in \mathbb{Z}[x]$ of ζ_{13} . Reinterpret your conjecture as a conjecture that relates the degrees of the irreducible factors of $f(x) \pmod{p}$ to the order of p modulo 13. Does your conjecture remind you of quadratic reciprocity?
5.
 - (a) Find by hand and with proof the ring of integers of each of the following two fields: $\mathbb{Q}(\sqrt{5})$, $\mathbb{Q}(i)$.
 - (b) Find the ring of integers of $\mathbb{Q}(x^5 + 7x + 1)$ using a computer.